

# North Atlantic Basin Modeling and Prediction for DAMEE-NAB

Robert C. Rhodes and Harley E. Hurlburt  
Ocean Dynamics and Prediction Branch  
Naval Research Laboratory  
Stennis Space Center, MS 39529-5004

phone: (228) 688-4704 (rhodes) fax: (228) 688-4759 email: [rhodes@nrlssc.navy.mil](mailto:rhodes@nrlssc.navy.mil)  
phone: (228) 688-4626 (hurlburt) fax: (228) 688-4759 email: [hurlburt@nrlssc.navy.mil](mailto:hurlburt@nrlssc.navy.mil)  
Award #: N0001498WX30188

<http://www7320.nrlssc.navy.mil/html/7320-home.html>

## LONG-TERM GOAL

An eddy-resolving nowcast/forecast system consisting of a  $1/8^\circ$  global system with embedded systems,  $1/32^\circ$  subtropical Atlantic ( $9-47^\circ\text{N}$ ) including the Caribbean and Gulf of Mexico and  $1/16^\circ$  Pacific north of  $20^\circ\text{S}$ .

## OBJECTIVES

The goal is an eddy-resolving prediction system for the North Atlantic Basin from  $9$  to  $47^\circ\text{N}$ , which includes the Caribbean and Gulf of Mexico. This project represents continued participation of NRL in the Data Assimilation and Model Evaluation Experiment – North Atlantic Basin (DAMEE-NAB) project supported by ONR with a focus on model-data comparisons and data assimilation. NRL has had two unique roles in DAMEE-NAB. The first was to investigate the impact of using extremely high horizontal resolution, the highest ever for basin-scale ocean modeling, because of the exceptional efficiency of the NRL Layered Ocean Model. The second was to assess different wind products as potential forcing for Atlantic Ocean models by investigating linear ocean model response to the different wind sets.

## APPROACH

The modeling effort is aimed at eddy-resolving models for the subtropical Atlantic Ocean and associated model development in collaboration with other projects. A wide variety of model-data comparisons are performed for evaluation of data assimilative and non-assimilative model experiments. Hydrodynamic and thermodynamic versions of NLOM are used with grid resolutions of  $1/4^\circ$  to  $1/64^\circ$  for each variable and 5 to 6 lagrangian layers in the vertical. The model has a free surface and allows diapycnal mixing, isopycnal outcropping and inflow/outflow through ports in the model boundaries. A version which includes a mixed layer and sea surface temperature is under development. The model runs efficiently and interchangeably on all DoD HPC platforms designed to handle applications this large, including massively parallel distributed memory computers and multi-processor shared memory computers. In general, NLOM is the most efficient ocean model in existence in terms of computer time per model year.

The altimeter heights will be assimilated using a combination of optimum interpolation, statistical inference, and nudging. However, the methods will depend upon the model chosen. Past experience with parameter optimization in a Gulf Stream regional system indicates that substantial improvements in

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>1998</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1998 to 00-00-1998</b>	
4. TITLE AND SUBTITLE <b>North Atlantic Basin Modeling and Prediction for DAMEE-NAB</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Research Laboratory, Ocean Dynamics and Prediction Branch, Stennis Space Center, MS, 39529</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM002252.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>8</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

performance can be obtained by empirically adjusting parameters used in the assimilation methods. More advanced methods of assimilation, such as the adjoint and Kalman filtering techniques, have not yet reached a state where they are practical alternatives.

Linear solutions are used for assessment of different wind products for potential ocean model forcing because they represent a deterministic oceanic response to the wind forcing and because they have well understood dynamics which represent the lowest order response to wind forcing, i.e. Sverdrup (1947) flow for the ocean gyre interiors which are closed by Munk (1950) western boundary layer currents.

## **WORK COMPLETED**

Two papers were submitted for a special issue of Deep-Sea Research devoted to DAMEE-NAB. The first investigates the impact of  $1/8^\circ$ ,  $1/16^\circ$ ,  $1/32^\circ$ , and  $1/64^\circ$  ocean model resolution on model-data comparisons for the Gulf Stream system between the Florida Straits and the Grand Banks. The  $1/64^\circ$  simulation (Fig. 1) is a first of a kind simulation, the highest horizontal resolution to date for a model of a major ocean basin. It was performed under an FY98 DoD High Performance Computing Challenge grant of computer time. Model-data comparisons included 1) the Gulf Stream mean pathway and its variability using GEOSAT-ERM and TOPEX/POSEIDON (T/P) altimetric analyses of Gulf Stream axis and IR analyses of the Gulf Stream north wall, 2) the mean position, eastward extent and amplitude of the Gulf Stream nonlinear recirculation gyres using surface dynamic height from the Navy's GDEM oceanic climatology, 3) mean transports of upper ocean and abyssal currents, especially the Gulf Stream and the Deep Western Boundary Current, 4) rms sea surface height variability from T/P (Fig. 2), and 5) abyssal eddy kinetic energy from current meter data. In addition, initial experiments were performed using a  $1/32^\circ$  thermodynamic version of the subtropical Atlantic model.

The second paper investigated linear Atlantic ocean model response to 11 different wind sets and a linear, Munk western boundary layer representation of the global thermohaline circulation. The wind stress climatologies used in this study were: 1) smoothed Hellerman and Rosenstein (HR, 1983), 2) unsmoothed HR, 3) Comprehensive Ocean-Atmosphere Data Set (COADS) (da Silva *et al.*, 1994a,b), 4) Isemer and Hasse (1987), 5) European Centre for Medium-Range Forecasts (ECMWF) 1000 mb from 1981-1993, 6) ECMWF 10 m from 1985-1994, 7) ECMWF 1000 mb from 1983-1986 (ECMWF, 1995), 8) ECMWF 1000 mb re-analysis from 1979-1993, 9) ECMWF 10 m re-analysis from 1979-1993 (Gibson *et al.*, 1997), 10) Fleet Numerical Meteorology and Oceanography Center (FNMOC) 7/90-6/95 (Hogan and Rosmond, 1991), and 11) National Centers for Environmental Prediction surface stress re-analysis for 1979-1995 (Kalnay *et al.*, 1996).


Atlantic region data assimilation was performed in the context of T/P altimetric data assimilation into a  $1/4^\circ$  global ocean model in collaboration with 6.2 project Basin Scale Ocean Prediction Systems and 6.4 project Ocean Data Assimilation. Particular attention was focussed on results obtainable from a  $1/4^\circ$  model with insufficient horizontal resolution for Gulf Stream separation and an inaccurate Gulf Stream mean pathway. Results from track by track assimilation were compared with assimilation of complete field OI analyses of SSH obtained using T/P data over a 10-day repeat interval.

## RESULTS

The paper about the impact of ocean model resolution ( $1/8^\circ$  to  $1/64^\circ$  for each variable) on model-data comparisons showed that while  $1/16^\circ$  resolution is a major step toward realistic simulations,  $1/32^\circ$  resolution is required for realistic and robust results. This includes Gulf Stream separation from the coast at Cape Hatteras, the Gulf Stream pathway between Cape Hatteras and the Grand Banks, the large-scale C-shape of the subtropical gyre, and the oceanic variability in the upper ocean and the abyssal layer.

The paper about the ocean model response to different wind forcing and a linear, Munk western boundary layer representation of the global thermohaline circulation showed that the latter makes a substantial contribution to the Gulf Stream transport and is essential for realistic circulation patterns in the Caribbean and western tropical Atlantic. Overall, a smoothed version of the Hellerman and Rosenstein (1983) and the COADS (da Silva, 1994a,b) wind stress climatologies, with the global thermohaline component added, gave the best results for the Gulf Stream region, including Florida Straits transport close to the well-observed value.

When assimilating satellite altimeter data into a  $1/4^\circ$  model with insufficient resolution for Gulf Stream separation and an inaccurate Gulf Stream mean pathway, complete field assimilation gives stronger constraint on the model and gives a better representation of the Gulf Stream, as well as improved results within the Gulf of Mexico. The weaker constraint of track by track assimilation allows the inaccurate  $1/4^\circ$  model Gulf Stream dynamics to “fight” the data too much. In principle, track by track assimilation is preferable because it is synoptic (whereas, the OI analyses are not) and because it uses the model as a dynamic interpolator.



**Figure 1.** Whole domain snapshot of SSH from a  $1/64^\circ$  5-layer Atlantic simulation forced by the Hellerman-Rosenstein monthly wind stress climatology and the global thermohaline circulation.

## IMPACT/APPLICATION

Simulation of realistic Gulf Stream separation from the coast at Cape Hatteras and a realistic Gulf Stream pathway between Cape Hatteras and the Grand Banks is a “Holy Grail” problem in ocean modeling, where many have tried and failed. Resolution requirements were assessed for realistic simulation of the Gulf Stream system and it was found that  $1/32^\circ$  resolution is required for robust realistic results. Resolution of  $1/16^\circ$  is the highest used outside NRL for Atlantic basin-scale ocean modeling. The linear Atlantic simulations can be used 1) to aid in the evaluation of wind sets for forcing ocean models and 2) to assist modellers in assessing and understanding the impact of nonlinearity and additional model physics and dynamical processes. The linear simulations provide a baseline for comparison which has well understood dynamics.

The plan is to embed a  $1/32^\circ$  Atlantic model into a  $1/8^\circ$  global model for transition as part of an operational global ocean nowcast/forecast system with transition to FNMOC by the end of FY01. This would include participation in the Global Ocean Data Assimilation Experiment (GODAE) as a pilot program, where the NRL effort is represented on both the international and U.S. Scientific Steering Teams. The system will include assimilation of real-time altimeter data received from NAVOCEANO’s


Altimeter Data Fusion Center. It would be an upgrade to an existing  $1/4^\circ$  global ocean model which already assimilates real-time altimeter data from T/P and ERS-2 (and soon probably from the Navy's GFO) and which is already running in real time at FNMOC.

## **TRANSITIONS**

A  $1/4^\circ$  6 layer finite depth thermodynamic global ocean model was transitioned to FNMOC in FY98 via the data assimilation group. This model is now running in real time with T/P and ERS-2 altimetry data assimilation. The system is being prepared for OPTTEST and is expected to be operational in FY99.

## **RELATED PROJECTS**

This 6.1 project and a collaborative 6.2 project, Basin-Scale Ocean Prediction Systems, are NRL participants in ONR's DAMEE-NAB project. The DAMEE-NAB project is multi-institutional. In addition to NRL, it includes Mississippi State University (Dave Dietrich and Avichal Mehra), MIT (Paola Rizzoli's group), Princeton University (Tal Ezer and George Mellor), Rutgers University (Scott Glenn and Dale Haidvogel's group), UCLA (Michael Ghil's group), U. Miami (Eric Chassignet's group) and the University of Southern Mississippi (Ziv Sirkes and Bob Willems). None of these participated directly in the NRL effort. In addition, this project has collaborated with other projects: 6.1 Dynamics of Low-Latitude Western Boundary Currents, 6.1 Forced Upper Ocean Dynamics, 6.1 Topographic and Thermodynamic forcing, 6.4 Ocean Data Assimilation, 6.4 Large-Scale Ocean Models and Data Assimilation in Ocean Prediction (in collaboration with J. Moura and A. Asif at Carnegie Mellon University).



*Figure 2. SSH variability (color) in the Gulf Stream region from (a) 1/16° 5-layer Atlantic simulation 16H, (b) 1/32° 5-layer Atlantic simulation 32H, (c) 1/64° 5-layer Atlantic simulation 64H, (d) 1/16° 6-layer global simulation forced by Hellerman-Rosenstein monthly climatological winds (simulation 16GH), (e) 1/16° 6-layer global simulation over 1990-1996 that was forced by interannual winds (simulation 16GI) and (f) T/P altimetry over the time period 1993-1996. All of the simulations also included constant global thermohaline forcing. Superimposed on each is the mean (center line), lines a standard deviation away from the mean on either side and extreme (outer lines) of the Gulf Stream axis as determined from T/P altimetry by Lee et al. (1997). The contour interval of the SSH variability is 2 cm.*

## REFERENCES

- da Silva, A.M., Young, C.C., and Levitus, S. (1994a) Atlas of surface marine data 1994. Volume 1: Algorithms and procedures. *NOAA Atlas NESDIS 6*, U.S. Dept. of Commerce, NOAA, NESDIS, 83 pp.
- da Silva, A.M., Young, C.C., and Levitus, S. (1994b) Atlas of surface marine data 1994. Volume 3: Anomalies of heat and momentum fluxes. *NOAA Atlas NESDIS 8*, U.S. Dept. of Commerce, NOAA, NESDIS, 413 pp.
- European Centre for Medium-Range Weather Forecasts (ECMWF) (1995) User guide to ECMWF products. *Meteorol. Bull. M3.2*, ECMWF, Reading, Berkshire, UK, 71 pp.
- Gibson, J.K., Kallberg, P., Uppala, S., Hernandez, A., Nomura, A. and Serrano, E. (1997) ECMWF Re-Analysis Project Report Series:1. ERA Description, ECMWF. Reading, Berkshire, UK, 71 pp.
- Hellerman, S. and Rosenstein, M. (1983) Normal monthly wind stress over the world ocean with error estimates. *J. Phys. Ocean.*, **13**, 1093-1104.
- Hogan, T.F. and Rosmond, T.E. (1991) The description of the Navy Operational Global Atmospheric Prediction System's spectral forecast model. *Mon. Wea. Rev.*, **119**, 1786-1815.
- Isemer, H.J., and Hasse, L. (1987) The Bunker climate atlas of the North Atlantic Ocean. Vol. 2 *Air-Sea Interactions*. Springer-Verlag, 256 pp.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K.C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D. (1996) The NCEP/NCAR 40-year Reanalysis Project. *Bull. Amer. Meteorol. Soc.*, **77**, 437-471.
- Lee, H., Glenn, S.M., and Haidvogel, D.B. (1997) A Gulf Stream synthetic geoid for the TOPEX altimeter. *J. Atmos. Ocean. Tech.*, (submitted).
- Munk, W.H. (1950) On the wind-driven ocean circulation. *J. Meteorol.*, **7**, 79-93.
- Sverdrup, H.U. (1947) Wind-driven currents in a baroclinic ocean: With application to the equatorial currents of the eastern Pacific. *Proc. Natl. Acad. Sci.*, **33**, 318-326.

## PUBLICATIONS

- Hogan, P.J. and Hurlburt, H.E. (1998) Dependence of Gulf Stream simulations on grid resolution,  $1/2^\circ$  to  $1/32^\circ$  degree. The EuroGOOS Atlantic Workshop Report, Le Provost, C. and Flemmin, N.C., eds., Southampton Oceanography Centre, UK, pp. 14-18.



Hogan, P.J. and Hurlburt, H.E. (1998) Realistic simulation of the Gulf Stream pathway from a  $1/32^\circ$  layered ocean model. *Challenges* (ARSC newsletter), Arctic Region Supercomputing Center, Fairbanks, AK, Vol. 6, No. 1, pp. 1, 5.

Hurlburt, H.E. and Hogan, P.J. (1998) Realistic simulation of the Gulf Stream pathway from a  $1/32^\circ$  layered ocean model. *High Performance Computing Contributions to DoD Mission Success 1998*, DoD High Performance Computing Modernization Office, pg. 109.

Hurlburt, H.E. and Hogan, P.J. (1998) Impact of  $1/8^\circ$  to  $1/64^\circ$  resolution on Gulf Stream model-data comparisons in basin-scale subtropical Atlantic Ocean models, *Deep-Sea Res.* (submitted).

Townsend, T.L., Hurlburt, H.E. and Hogan, P.J. (1998) Modeled Sverdrup flow in the North Atlantic from eleven different wind stress climatologies, *Deep-Sea Res.* (submitted).

---